

NASA-CR-172,573

NASA Contractor Report 172573

ICASE REPORT NO. 85-17

NASA-CR-172573
19850017917

ICASE

FOR REFERENCE

NOT TO BE TAKEN FROM THIS LOCATION

ANALYSIS OF SPECTRAL OPERATORS IN ONE-DIMENSIONAL DOMAINS

Yvon Maday

Contract No. NAS1-17070

February 1985

INSTITUTE FOR COMPUTER APPLICATIONS IN SCIENCE AND ENGINEERING
NASA Langley Research Center, Hampton, Virginia 23665

Operated by the Universities Space Research Association

NASA

National Aeronautics and
Space Administration

Langley Research Center
Hampton, Virginia 23665

LIBRARY COPY

1985

LANGLEY RESEARCH CENTER
LIBRARY, NASA
HAMPTON, VIRGINIA

ANALYSIS OF SPECTRAL OPERATORS IN ONE-DIMENSIONAL DOMAINS

Yvon Maday

Laboratoire d'Analyse Numérique

Université Pierre et Marie Curie

and

Université de Paris Val de Marne

ABSTRACT

We prove results concerning certain projection operators on the space of all polynomials of degree less than or equal to N with respect to a class of one-dimensional weighted Sobolev spaces. These results are useful in the theory of the approximation of partial differential equations with spectral methods.

Research was supported in part by the National Aeronautics and Space Administration under NASA Contract No. NAS1-17070 while the author was in residence at the Institute for Computer Applications in Science and Engineering, NASA Langley Research Center, Hampton, VA 23665.

I. INTRODUCTION

This paper presents an investigation of a class of projection operators that arises in the analysis of the approximation of differential equations by spectral methods using Chebyshev decomposition.

Some similar operators have been studied before by Canuto-Quarteroni [1] and Maday-Quarteroni [1], but the existing results are not adequate in many applications. In fact they forbid analysis for the error of the approximation by spectral methods of fourth-order problems and, in several instances, second-order problems (see Canuto-Quarteroni [2]).

We first present some background tools required for our analysis. They consist of Sobolev spaces relative to the weight $\omega(x) = (1 - x^2)^{-1/2}$ (this weight arises in the relations of orthogonality of Chebyshev polynomials). We recall and complete results proved by Grisvard [1], [2] concerning interpolation theory between these spaces.

Then we present an analysis of projection operators from these spaces into the set of all polynomials of degree lower than N .

Finally we give an application of the results herein proved to a simple test problem.

We shall give other applications in Maday [1] and shall in a future work extend these results to multidimensional domains. Our aim is to apply such results to the analysis of the approximation of Navier-Stokes equations by spectral methods (see Maday-Metivet [1], [2]).

For some different notions about projection operators that arise in spectral methods, see Tadmor [1].

II. PRELIMINARIES: SOME FUNCTION SPACES

Notations and Basic Properties

Let J be an open interval $]a, b[$ of \mathbb{R} ($a < b$); we consider a weight function $\rho(x)$, continuous over J , satisfying $\rho(x) \geq \rho_0 > 0$ for any $x \in J$.

Let us set:

$$(2.1) \quad L^2_\rho(J) = \{ \phi : J \rightarrow \mathbb{R} \mid \phi \text{ is measurable and } (\phi, \phi)_\rho < +\infty \},$$

equipped with the inner product $(\phi, \psi)_\rho = \int_J \phi(x)\psi(x)\rho(x)dx$. For any integer $s \geq 0$ we set:

$$H^s_\rho(J) = \{ \phi \in L^2_\rho(J) \mid \|\phi\|_{s,\rho} < \infty \},$$

where:

$$(2.2) \quad \|\phi\|_{s,\rho} = ((\phi, \phi))_{s,\rho},$$

this space being equipped with the inner product:

$$(2.3) \quad ((\phi, \psi))_{s,\rho} = \sum_{k=0}^s \left(\frac{d^k \phi}{dx^k}, \frac{d^k \psi}{dx^k} \right)_\rho.$$

Clearly, one has the equality:

$$L^2_\rho(I) = H^0_\rho(I).$$

For any real $s \geq 0$, noninteger, $H^s_\rho(J)$ is defined by interpolation between the space $H^{\bar{s}}_\rho(J)$ and $H^{\bar{s}+1}_\rho(J)$, where \bar{s} represents the integral part of s .

The method of interpolation can be the complex one, the operator's domain one or the trace one (see Lions-Magenes [1] for more details). Besides we define $H_{0,\rho}^s(J)$ as being the closure of $\mathcal{D}(J)$ in $H_\rho^s(J)$. When $\rho = 1$ these spaces are the usual Sobolev spaces denoted by $H^s(J)$ and $H_0^s(J)$ respectively. For the application to spectral methods we are mostly interested in those spaces when $J = I \equiv]-1, +1[$ and $\rho(x) = \omega(x) \equiv \frac{1}{\sqrt{1-x^2}}$. Let us recall some results proved in Grisvard [1], [2] valuable for $J = I$, $\rho = \omega$ and for $J =]0, 1[, \rho = \frac{1}{\sqrt{x}}$.

THEOREM 2.1 (Grisvard [1]):

i) For any real $s > 0$, $s \notin \mathbb{N} + 1/4$ we have:

$$(2.4) \quad H_{0,\rho}^s(J) = [H_{0,\rho}^{\bar{s}}(J), H_{0,\rho}^{\bar{s}+1}(J)]_{[s-\bar{s}]}$$

ii) For any integral n we have:

$$(2.4') \quad [H_{0,\rho}^n(J), H_{0,\rho}^{n+1}(J)]_{1/4} \subset H_{0,\rho}^{n+1/4}(J).$$

iii) For any real $s > 0$, $s \notin \mathbb{N} + 1/2$:

$$H_\rho^s(J) \subset C^m(\bar{J}),$$

the space of continuous mapping defined over \bar{J} whose derivative of order $\leq m$ are continuous over \bar{J} , with $m = \overline{s - \frac{1}{2}}$.

The trace application defined from $C^\infty(\bar{J})$ into \mathbb{R}^{2n} :

$$u \longmapsto \left(u(-1), \frac{du}{dx}(-1), \dots, \frac{d^n u}{dx^n}(-1), u(1), \frac{du}{dx}(1), \dots, \frac{d^n u}{dx^n}(1) \right)$$

can be extended to a continuous mapping from $H_{\rho}^{n+1/4+\varepsilon}(I)$ onto \mathbb{R}^{2n} for any $\varepsilon > 0$.

iv) For any real $1/4 < s < 5/4$, $H_{0,\rho}^s(J)$ coincide with the subspace of $H_{\rho}^s(J)$ of functions vanishing at the real boundaries of J .

v) For any real $s > 1/2$, $H_{\rho}^s(J)$ is an algebra.

THEOREM 2.2 (Grisvard [2]):

For any $0 \leq q < s < p$, $H_{\rho}^s(J)$ satisfies the following double topological imbedding:

$$(2.5) \quad [H_{\rho}^q(J), H_{\rho}^p(J)]_{\theta,1} \subset H_{\rho}^s(J) \subset [H_{\rho}^q(J), H_{\rho}^p(J)]_{\theta,\infty},$$

with $\theta = \frac{s-p}{q-p}$, and the notation holds for the real interpolation (see Lions and Peetre [1]).

The two following results can be found in Canuto-Quarteroni [2] and Maday-Quarteroni [1].

THEOREM 2.3:

- i) For any real $s \geq 1/4$, $H^s(I) \subset H_{\omega}^{s-1/4}(I)$.
- ii) For any $0 \leq r < s$, the imbedding $H_{\omega}^s(I) \subset H_{\omega}^r(I)$ is compact.

In the next section we shall generalize the results (2.4) and (2.4').

Some New Results About Interpolation Between $H_{0,\omega}^p(I)$

This section is devoted to the proof of the following:

THEOREM 2.4: For any $0 \leq q < s < p$ not in $\mathbb{N} + 1/4$ we have:

$$[H_{0,\omega}^q(I), H_{0,\omega}^p(I)]_{\left[\frac{s-p}{q-p}\right]} = H_{0,\omega}^s(I).$$

This theorem is a consequence of the two following lemma:

LEMMA 2.1: For any integer $p \leq n$, we have:

$$u \in H_{0,\omega}^n(I) \Rightarrow \frac{d^p u}{dx^p} \in L_{\omega}^2{}_{4(n-p)+1}(I).$$

PROOF: It is an easy matter to check that this result is a consequence of

$$(2.6) \quad u \in H_{0, \frac{1}{\sqrt{x}}}^n(0,1) \Rightarrow \frac{d^p u}{dx^p} \in L_{\left(\frac{1}{\sqrt{x}}\right)^{4(n-p)+1}}^2(0,1),$$

(we shift the difficulties at ± 1 onto 0). So let u be in $H_{0, \frac{1}{\sqrt{x}}}^n(0,1)$; from Theorem 2.1 (point iv) we have, for any $0 \leq p < n$:

$$\frac{d^p u}{dx^p}(0) = 0,$$

hence

$$\int_0^x \frac{d^{p+1} u}{dx^{p+1}}(t) dt = \frac{d^p u}{dx^p}(x).$$

Besides, from Lemma 6.2.1 of Nečas [1] we have, for any $\alpha < 1$ and any v such that $\int_0^1 v^2(x) x^\alpha dx < \infty$:

$$(2.7) \quad \int_0^1 \left(\int_0^x |v(x)| \right)^2 x^{\alpha-2} dx \leq \left(\frac{1}{1-\alpha} \right) \int_0^1 |v(x)|^2 x^\alpha dx,$$

taking then $\alpha = -\frac{1}{2} - 2(n - (p+1))$ and $v = \frac{d^{p+1}u}{dx^{p+1}}$ we obtain:

$$\int_0^1 \left(\frac{d^p u}{dx^p} \right)^2 x^{-1/2 - 2(n-p)} dx \leq C \int_0^1 \left(\frac{d^{p+1}u}{dx^{p+1}} \right)^2 x^{-1/2 - 2(n-(p+1))} dx,$$

and (2.6) holds by induction over p .

LEMMA 2.2: For any integer $n > 0$, the mapping $u \mapsto u\omega^{1/2}$ is an homeomorphism from $H_{0,\omega}^n(I)$ onto $H_0^n(I)$.

PROOF: Here again we prove the result for the weight $\frac{1}{\sqrt{x}}$, say:

$$(2.8) \quad u \mapsto ux^{-1/4} \text{ is an homeomorphism from } H_{0,\frac{1}{\sqrt{x}}}^n(0,1) \text{ onto } H_0^n(0,1).$$

Let $\phi \in \mathcal{D}(0,1)$, then, for $0 \leq m \leq n$:

$$\begin{aligned} \frac{d^m}{dx^m}(\phi x^{-1/4}) &= \sum_{p=0}^m C_m^p \frac{d^p \phi}{dx^p} \frac{d^{m-p}(x^{-1/4})}{dx^{m-p}} \\ &= \sum_{p=0}^m C_m^p D_m^p \frac{d^p \phi}{dx^p} x^{-1/4 - (m-p)}, \end{aligned}$$

with $D_m^p = [-1/4 - (m-p+1)]D_m^{p+1}$ and $D_m^m = 1$. From Lemma 2.1 we then get:

$$\left\| \frac{d^m}{dx^m} (\phi x^{-1/4}) \right\|_{0,1} \leq C \|\phi\|_{m,1/\sqrt{x}} \leq C \|\phi\|_{n,1/\sqrt{x}} ;$$

summing up these estimates for $0 \leq m \leq n$ we derive:

$$(2.9) \quad \|\phi x^{-1/4}\|_{n,1} \leq C \|\phi\|_{n,1/\sqrt{x}} .$$

Inversely, let us prove that, for any $\phi \in \mathcal{D}(I)$:

$$(2.10) \quad \|\phi x^{1/4}\|_{n,1/\sqrt{x}} \leq C \|\phi\|_{n,1} .$$

From Hardy's inequality (Lemma 2.5.1 of Nečas [1]) we derive by induction that, for any $0 \leq p \leq m \leq n$:

$$(2.11) \quad \left\| \frac{d^p \phi}{dx^p} \right\|_{0,x^{-2(m-p)}} \leq C \|\phi\|_{m,1} ,$$

besides:

$$\begin{aligned} \frac{d^m (\phi x^{1/4})}{dx^m} &= \sum_{p=0}^m C_m^p \frac{d^p \phi}{dx^p} \frac{d^{m-p}}{dx^{m-p}} (x^{1/4}) \\ &= \sum_{p=0}^m C_m^p D_m^p \frac{d^p \phi}{dx^p} x^{1/4 - (m-p)} , \end{aligned}$$

with:

$$D_m^p = [1/4 - (m - p + 1)] D_m^{p+1} \quad \text{and} \quad D_m^m = 1 . .$$

Then using (2.11) we get:

$$\frac{d^m(\phi x^{1/4})}{dx^m} \in L^2_{1/\sqrt{x}}(0,1),$$

and (2.10) is derived by summing up these results for $0 \leq m \leq n$. We can now achieve (2.8) as a consequence of (2.9) and (2.10).

We can now prove the main result of this section.

PROOF OF THEOREM 2.4: From (2.4) and Lemma 2.2 we deduce that the mapping $u \mapsto u\omega^{1/2}$ is an homeomorphism from $H^s_{0,\omega}(I)$ onto $H^s_0(I)$ for any $s \geq 0$ not in $\mathbb{N} + 1/4 \cap \mathbb{N} + 1/2$ (see Lions-Magenes [1] for more details about the properties of spaces of interpolation).

Let us recall that, for any $q \leq s \leq p$ not in $\mathbb{N} + 1/2$ we have (see Lions-Magenes [1]):

$$(2.12) \quad H^s_0(I) = [H^q_0(I), H^p_0(I)]_{\frac{s-q}{p-q}}.$$

From the previous homeomorphism we deduce that, for any $q \leq s \leq p$ not in $\{\mathbb{N} + 1/2\} \cup \{\mathbb{N} + 1/4\}$:

$$(2.13) \quad H^s_{0,\omega}(I) = [H^q_{0,\omega}(I), H^p_{0,\omega}(I)]_{\frac{s-q}{p-q}}.$$

Let us remark now that the values of p, q, s in $\mathbb{N} + 1/2$ have only been excluded due to (2.12), these values can now be recovered thanks to the reiteration theorem (Theorem I.6.1 of Lions-Magenes [1]).

III. APPROXIMATION RESULTS OF PROJECTION OPERATOR IN WEIGHTED SOBOLEV SPACES

The previous theorem leads us to define over $H_{0,\omega}^r(I)$ a new scalar product. Indeed, for p not in $\mathbb{N} + 1/4 \cup \mathbb{N} + 1/8$, $H_{0,\omega}^p(I)$ can be seen as the interpolate $1/2$ between $L_{\omega}^2(I)$ and $H_{0,\omega}^{2p}(I)$ and for p in $\mathbb{N} + 1/8$, $H_{0,\omega}^p(I)$ can be seen as the interpolate $1/3$ between $L_{\omega}^2(I)$ and $H_{0,\omega}^{3p}(I)$.

If we consider the domain operator interpolation, this find expression in the existence of a selfadjoint operator Λ_r such that:

- * if $r \in \mathbb{N} + 1/8$, the domain $D(\Lambda_r^3)$ of the operator Λ_r^3 in $L_{\omega}^2(I)$ is $H_{0,\omega}^{3r}(I)$ if $r \notin \mathbb{N} + 1/8$, the domain of $D(\Lambda_r^2)$ of the operator Λ_r^2 in $L_{\omega}^2(I)$ is $H_{0,\omega}^{2r}(I)$.
- * The domain $D(\Lambda_r)$ of the operator Λ_r in $L_{\omega}^2(I)$ is $H_{0,\omega}^r(I)$ if $r \notin \mathbb{N} + 1/4$ and is included in $H_{0,\omega}^r(I)$ if $r \in \mathbb{N} + 1/4$.

Moreover:

$$(3.1) \quad (u, v) \longmapsto (((u, v)))_{r, \omega} \equiv (\Lambda_r u, \Lambda_r v)_{\omega},$$

is a scalar product whose associated norm is equivalent to the one defined in (2.2) if $r \notin \mathbb{N} + 1/4$.

Let us define now $P_{r,N}$ as the projection operator from $H_{0,\omega}^r(I)$ over S_N^r with respect to the previous scalar product with:

$$S_N^r = S_N \cap H_{0,\omega}^r(I),$$

$$S_N = \{ \phi \text{ defined over } I \mid \phi \text{ is a polynomial of degree } \leq N \}.$$

LEMMA 3.1: Let $0 \leq v \leq r \leq \sigma$ with $\sigma \notin \mathbb{N} + 1/4$ we have, for any $\phi \in H_{\omega}^{\sigma}(I) \cap H_{0,\omega}^r(I)$:

$$(3.2) \quad \|\phi - P_{r,N} \phi\|_{v,\omega} \leq C N^{v-\sigma} \|\phi\|_{\sigma,\omega}.$$

REMARK 3.1: The case $v = r = 0$ has been studied in Canuto-Quarteroni [1], the case $0 \leq v \leq r = 1$ has been looked at in Maday-Quarteroni [1] (note that the dependence of the constant is then $C(\sigma) = C \cdot (\sigma!)$). Moreover it is proved that no optimal bound was possible for $H_{0,\omega}^v(I)$ norms with $v > r$. Indeed, for example:

$$(3.4) \quad \|\phi - P_{0,N} \phi\|_{v,\omega} \leq C N^{2v-\sigma} \|\phi\|_{\sigma,\omega}.$$

It is often necessary (see Canuto-Quarteroni [2], Maday-Metivet [2], Maday [1], and (4.14)) to obtain optimal results in higher norms.

PROOF: We shall only consider the case $r \notin \mathbb{N} + 1/8$ for simplicity. The proof is divided in two stages

i) We first prove (3.2) by induction over r in \mathbb{N} . So, let us assume that (3.2) is true for $s < r$ in \mathbb{N} ; let $\phi \in H_{0,\omega}^r(I)$; then $\phi_x \in H_{0,\omega}^{r-1}(I)$ and $P_{r-1,N-1}(\phi_x) \in S_{N-1}^{r-1}$. Moreover if $\phi(-1) = \phi(1) = 0$ we have:

$$\alpha \equiv \int_{-1}^1 P_{r-1,N-1}(\phi_x)(t) dt = \int_{-1}^1 [P_{r-1,N-1}(\phi_x) - \phi_x](t) dt.$$

From the Cauchy-Schwarz inequality we derive:

$$|\alpha| \leq \left(\int_{-1}^1 (P_{r-1, N-1}(\phi_x) - \phi_x)^2(t) \omega(t) dt \right)^{1/2} \left(\int_{-1}^1 (\omega(t))^{-1} dt \right)^{1/2}$$

$$\leq C \|P_{r-1, N-1}(\phi_x) - \phi_x\|_{0, \omega};$$

hence, from the induction hypothesis:

$$(3.5) \quad |\alpha| \leq C N^{1-\sigma} \|\phi_x\|_{\sigma-1, \omega}.$$

Finally we have:

$$R_N(x) = \int_{-1}^x \left[P_{r-1, N-1}(\phi_x)(t) - \frac{\alpha(1-t^2)^{r-1}}{\int_{-1}^1 (1-x^2)^{r-1} dx} \right] dt \in S_N^r.$$

Due to the Poincaré-like inequality, the polynomial satisfies the following:

$$\|\phi - R_N\|_{r, \omega} \leq \|(\phi - R_N)_x\|_{r-1, \omega},$$

the induction hypothesis, and (3.5) gives us:

$$\|\phi - R_N\|_{r, \omega} \leq C(N^{(r-1)-(\sigma-1)} + N^{(1-\sigma)}) \|\phi_x\|_{\sigma-1, \omega}$$

$$\leq C N^{r-\sigma} \|\phi\|_{\sigma, \omega}.$$

From the equivalence of the norms $\|\cdot\|_{r, \omega}$ and $|||\cdot|||_{r, \omega}$, and the identity:

$$|||\phi - P_{r, N} \phi|||_{r, \omega} = \inf_{\phi_N \in S_N^r} |||\phi - \phi_N|||_{r, \omega},$$

we obtain for any ϕ in $H_{\omega}^{\sigma}(I) \cap H_{0,\omega}^r(I)$:

$$(3.6) \quad |||\phi - P_{r,N} \phi|||_{r,\omega} \leq CN^{r-\sigma} \|\phi\|_{\sigma,\omega}.$$

Besides, since the operator Λ_r is selfadjoint, we have:

$$\begin{aligned} \|\phi - P_{r,N} \phi\|_{0,\omega} &= \inf_{\psi \in L_{\omega}^2(I)} \frac{(\phi - P_{r,N} \phi, \psi)_{\omega}}{\|\psi\|_{0,\omega}} \\ &= \inf_{\psi \in L_{\omega}^2(I)} \frac{(\Lambda_r(\phi - P_{r,N} \phi), \Lambda_r^{-1} \psi)_{\omega}}{\|\psi\|_{0,\omega}}. \end{aligned}$$

From (3.1) we then get:

$$(3.7) \quad \|\phi - P_{r,N} \phi\|_{0,\omega} = \inf_{\psi \in L_{\omega}^2(I)} \frac{(((\phi - P_{r,N} \phi, \Lambda_r^{-2} \psi)))_{r,\omega}}{\|\psi\|_{0,\omega}}.$$

By definition of $P_{r,N}$ we have, for any ψ in $L_{\omega}^2(I)$:

$$(((\phi - P_{r,N} \phi, P_{r,N}(\Lambda_r^{-2} \psi))))_{r,\omega} = 0;$$

hence

$$\begin{aligned} \|\phi - P_{r,N} \phi\|_{0,\omega} &= \inf_{\psi \in L_{\omega}^2(I)} \frac{(((\phi - P_{r,N} \phi, (\Lambda_r^{-2} \psi) - P_{r,N}(\Lambda_r^{-2} \psi))))_{r,\omega}}{\|\psi\|_{0,\omega}} \\ &\leq |||\phi - P_{r,N} \phi|||_{r,\omega} \inf_{\psi \in L_{\omega}^2(I)} \frac{|||(\Lambda_r^{-2} \psi) - P_{r,N}(\Lambda_r^{-2} \psi)|||_{r,\omega}}{\|\psi\|_{0,\omega}}. \end{aligned}$$

Due to (3.6) we then derive:

$$\begin{aligned}
 \|\phi - P_{r,N} \phi\|_{0,\omega} &\leq CN^{r-\sigma} \|\phi\|_{\sigma,\omega} N^{-r} \inf_{\psi \in L_{\omega}^2(I)} \frac{\|\Lambda_r^{-2} \psi\|_{0,\omega}}{\|\psi\|_{0,\omega}} 2r, \omega \\
 &\leq CN^{-\sigma} \|\phi\|_{\sigma,\omega} \inf_{\psi \in L_{\omega}^2(I)} \frac{\|\Lambda_r^2(\Lambda_r^{-2} \psi)\|_{0,\omega}}{\|\psi\|_{0,\omega}} \\
 &\leq CN^{-\sigma} \|\phi\|_{\sigma,\omega}.
 \end{aligned}$$

Now, from the two estimates, valuable for any $\phi \in H_{\omega}^{\sigma}(I) \cap H_{0,\omega}^r(I)$:

$$\|\phi - P_{r,N} \phi\|_{r,\omega} \leq CN^{r-\sigma} \|\phi\|_{\sigma,\omega}$$

$$\|\phi - P_{r,N} \phi\|_{0,\omega} \leq CN^{-\sigma} \|\phi\|_{\sigma,\omega}$$

we derive that for any $\theta \in]0,1[$:

$$\|\phi - P_{r,N} \phi\|_{[L_{\omega}^2(I), H_{0,\omega}^r(I)]_{\theta}} \leq CN^{\theta r - \sigma} \|\phi\|_{\sigma,\omega}.$$

Due to (2.4) and (2.4') we deduce that, for any $0 \leq \nu \leq r$:

$$(3.8) \quad \|\phi - P_{r,N} \phi\|_{\nu,\omega} \leq CN^{\nu - \sigma} \|\phi\|_{\sigma,\omega}.$$

ii) Let us now prove (3.3) for nonintegral values of r . Let $\phi \in \mathcal{D}(I)$, from step (i) we know that, for any $\sigma \geq \overline{r+1}$, $\sigma \notin \mathbb{N} + 1/4$

$$\inf_{\phi_N \in S_N^{\overline{r+1}}} \|\phi - \phi_N\|_{\overline{r+1}, \omega} \leq C N^{\overline{r+1}-\sigma} \|\phi\|_{\sigma, \omega},$$

$$\inf_{\phi_N \in S_N^{\overline{r+1}}} \|\phi - \phi_N\|_{\overline{r}, \omega} \leq C N^{\overline{r}-\sigma} \|\phi\|_{\sigma, \omega}$$

(see (3.8) with $v = \overline{r+1}$ and $v = \overline{r}$ respectively). These two estimates are equivalent to the following one:

$$\|\dot{\phi}\|_{H_{0,\omega}^{\overline{r+1}}(I)/S_N^{\overline{r+1}}} \leq C N^{\overline{r+1}-\sigma} \|\phi\|_{\sigma, \omega},$$

$$\|\dot{\phi}\|_{H_{0,\omega}^{\overline{r}}(I)/S_N^{\overline{r+1}}} \leq C N^{\overline{r}-\sigma} \|\phi\|_{\sigma, \omega}.$$

Due to the interpolation of quotient spaces (see Lions-Magenes [1] Lemma I.13.2) we have, for any $\theta \in]0,1[$:

$$\|\dot{\phi}\|_{[H_{0,\omega}^{\overline{r}}(I), H_{0,\omega}^{\overline{r+1}}(I)]_{\theta}/S_N^{\overline{r+1}}} \leq C N^{\overline{r}+\theta-\sigma} \|\phi\|_{\sigma, \omega}.$$

From (2.4), (2.4') we deduce (we take $\theta = r - \overline{r}$):

$$\|\dot{\phi}\|_{H_{0,\omega}^r(I)/S_N^{\overline{r+1}}} \leq C N^{r-\sigma} \|\phi\|_{\sigma, \omega},$$

so that, for any $\sigma \geq \overline{r+1}$ $\sigma \notin \mathbb{N} + 1/4$:

$$(3.9) \quad \|\phi - P_{r,N} \phi\|_{r,\omega} \leq CN^{r-\sigma} \|\phi\|_{\sigma, \omega}.$$

By definition we know that, for any $r \notin \mathbb{N} + 1/4$.

$$\|\phi - P_{r,N} \phi\|_{r,\omega} \leq C \|\phi\|_{r,\omega};$$

hence, (3.9) holds for any $\sigma \geq r$, $\sigma \notin \mathbb{N} + 1/4$ and any $r \in \mathbb{R}^{+*}$. The estimates in lower order norms are obtained following the same lines as in i).

We are now interested in the approximation of the spaces $H_{\omega}^r(I) \cap H_{0,\omega}^s(I)$ for $s \in \mathbb{N}$, $0 \leq s \leq r$ by polynomials therein contained.

For simplicity of exposition we shall consider the case $s = 0$. From point iii) of Theorem 2.4 we can easily exhibit for any $\phi \in H_{\omega}^r(I)$ a polynomial ϕ_0 of degree $\leq 2r - 1$ such that: $\phi - \phi_0 \in H_{0,\omega}^r(I)$ and for any real p :

$$(3.10) \quad \|\phi_0\|_{p,\omega} \leq C \|\phi\|_{r,\omega}.$$

Due to the previous lemma, we have, for any $0 \leq v \leq r \leq \sigma \notin \mathbb{N} + 1/4$:

$$\|(\phi - \phi_0) - (P_{r,N}(\phi - \phi_0))\|_{v,\omega} \leq CN^{v-\sigma} \|\phi - \phi_0\|_{\sigma, \omega},$$

and (3.10) then implies:

$$\|\phi - (\phi_0 + P_{r,N}(\phi - \phi_0))\|_{v,\omega} \leq CN^{v-\sigma} \|\phi\|_{\sigma, \omega}.$$

For N large enough (more precisely $N \geq 2r - 1$) we then get the existence of an operator \tilde{P} from $H_{\omega}^r(I)$ onto S_N such that:

$$\|\phi - \tilde{P}(\phi)\|_{v,\omega} \leq CN^{v-\sigma} \|\phi\|_{\sigma,\omega}.$$

This estimate provides an answer to our question in the case $s = 0$. The same proof can be done to build an operator from $H_{\omega}^r(I) \cap H_{0,\omega}^s(I)$ onto S_N^s satisfying analogous bounds. This leads us to state the main theorem of this paper:

THEOREM 3.1: Let $(v,r,\sigma) \in \mathbb{R}$, $\sigma \notin \mathbb{N} + 1/4$, $s \in \mathbb{N}$, $0 \leq s \leq r$, $0 \leq v \leq r \leq \sigma$. There exists an operator $\Pi_{r,N}^{s,0}$ from $H_{\omega}^r(I) \cap H_{0,\omega}^s(I)$ onto S_N^s such that, for any $\phi \in H_{\omega}^{\sigma}(I) \cap H_{0,\omega}^s(I)$ we have:

$$\|\phi - \Pi_{r,N}^{s,0} \phi\|_{v,\omega} \leq CN^{v-\sigma} \|\phi\|_{\sigma,\omega}.$$

IV. AN APPLICATION

Definition of the Problem

In order to explain how the previous results can be applied, we shall study an approximation of the very simple problem:

Find ψ defined over I such that:

$$(4.1) \quad \begin{cases} \frac{d^4 \psi}{dx^4} = f & \text{over } I, \\ \psi = \frac{d\psi}{dx} = 0 & \text{at } \pm 1. \end{cases}$$

(This problem provides a first step for the analysis of Stokes and Navier-Stokes problems in the ψ -formulation; see Maday-Metivet [1], [2].) Let us define $H_{\omega}^{-2}(I)$ as follows:

$$H_{\omega}^{-2}(I) = \{f \in \mathcal{D}'(I) \mid \exists g \in L_{\omega}^2(I): f = \frac{d^2 g}{dx^2}\}.$$

We now want to prove the following:

THEOREM 4.1: Let $f \in H_{\omega}^{-2}(I)$; then there exists one and only one solution ψ to the problem (4.1) in the space $H_{0,\omega}^2(I)$.

This theorem is a very simple consequence of Lax Milgram lemma and the two following lemmas.

LEMMA 4.1: There exist two positive constants δ_1 and δ_2 such that, for any ϕ in $H_{0,\omega}^2(I)$:

$$\int_I \phi^2 \omega^9 \leq \delta_1 \int_I \frac{d\phi^2}{dx} \omega^5 \leq \delta_2 \int_I \left(\frac{d^2 \phi}{dx^2}\right)^2 \omega.$$

This lemma is a corollary of Lemma 2.1.

LEMMA 4.2: There exist 3 positive constants α, β, γ such that for any $(\phi, \psi) \in H_{\omega}^2(I) \times H_{0, \omega}^2(I)$:

$$(4.2) \quad (\psi_{xx}, (\psi\omega)_{xx}) \geq \alpha \|\psi\|_{2, \omega}^2,$$

$$(4.3) \quad \|\psi\|_{2, \omega} \leq \beta \|\psi_{xx}\|_{0, \omega},$$

$$(4.4) \quad (\phi_{xx}, (\psi\omega)_{xx})_{\omega} \leq \gamma \|\phi_{xx}\|_{0, \omega} \|\psi_{xx}\|_{0, \omega}.$$

PROOF:

i) We first note that (4.3) is an easy consequence of the previous lemma, and is equivalent to the Poincaré inequality.

ii) Next, we get the following equalities, for any $\psi \in \mathcal{D}(I)$:

$$\begin{aligned} \int_I \psi_{xx} (\psi\omega)_{xx} dx &= \int_I \psi_{xx}^2 \omega dx + 2 \int_I \psi_{xx} \psi_x \omega_x dx + \int_I \psi_{xx} \psi \omega_{xx} dx \\ &= \int_I \psi_{xx}^2 \omega dx + \int_I (\psi_x^2)_x \omega_x dx + \int_I (\psi_x)_x \psi \omega_{xx} dx \\ &= \int_I \psi_{xx}^2 \omega dx - \int_I \psi_x^2 \omega_{xx} dx - \int_I \psi_x (\psi \omega_{xx})_x dx \\ &= \int_I \psi_{xx}^2 \omega dx - 2 \int_I \psi_x^2 \omega_{xx} dx + 1/2 \int_I \psi^2 \omega_{xxxx} dx \end{aligned}$$

let us note that:

$$\omega_{xx} = (1 + 2x^2)\omega^5,$$

$$\omega_{xxxx} = (9 + 72x^2 + 24x^4)\omega^9;$$

hence:

(4.5)

$$\int_I \psi_{xx}(\psi\omega)_{xx} dx = \int_I \psi_{xx}^2 dx - 2 \int_I \psi_x^2 (1 + 2x^2)\omega^5 dx + 1/2 \int_I \psi^2 (9 + 72x^2 + 24x^4)\omega^9 dx.$$

Besides, let us set:

$$P \equiv \int_I (\psi_{xx} \omega + 2x\psi_x \omega^3 + (2x^2 + 10^{-2})\psi\omega^5)^2 \omega^{-1} dx;$$

P is ≥ 0 and, an easy calculation gives:

$$P \equiv \int_I \psi_{xx}(\psi\omega)_{xx} - 2.10^{-2} \int_I \psi_x^2 \omega^5 - \int_I (5.78x^2 + 0.4839)\psi^2 \omega^9,$$

so that

$$\int_I \psi_x^2 \omega^5 \leq 50 \int_I \psi_{xx}(\psi\omega)_{xx},$$

from which we derive:

$$\int_I \psi_x^2 (1 + 2x^2)\omega^5 \leq 150 \int_I \psi_{xx}(\psi\omega)_{xx}.$$

Using that inequality in (4.5) we obtain $\alpha > 0$ such that:

$$\int_I \psi_{xx} (\psi \omega)_{xx} \geq \alpha \beta \int_I \psi_{xx}^2 \omega,$$

and (4.2) is a consequence of (4.3).

iii) Finally, let us note that for any (ϕ, ψ) in $H_\omega^2(I) \times H_{0,\omega}^2(I)$ we have:

$$(4.6) \quad (\phi_{xx}, (\psi \omega)_{xx})_1 = (\phi_{xx}, \psi_{xx})_\omega + 2 \int_I \phi_{xx} \psi_x \omega_x + \int_I \phi_{xx} \psi \omega_{xx}.$$

The following inequality is simple:

$$(4.7) \quad |(\phi_{xx}, \psi_{xx})_\omega| \leq \|\phi_{xx}\|_{0,\omega} \|\psi_{xx}\|_{0,\omega}.$$

Let us examine the second term:

$$\left| \int_I \phi_{xx} \psi_x \omega_x \right| = \left| \int_I \phi_{xx} (\psi_x \omega_x \omega^{-1}) \omega \right| \leq \left| \int_I \phi_{xx}^2 \omega \right|^{1/2} \left| \int_I \psi_x^2 \omega_x^2 \omega^{-1} \right|^{1/2};$$

since $\omega_x^2 \omega^{-1} = x^2 \omega^5$, we derive from Lemma 4.1 that:

$$(4.8) \quad \left| \int_I \phi_{xx} \psi_x \omega_x \right| \leq C \|\phi_{xx}\|_{0,\omega} \|\psi_{xx}\|_{0,\omega}.$$

We obtain, in a similar way:

$$(4.9) \quad \left| \int_I \phi_{xx} \psi \omega_{xx} \right| \leq C \|\phi_{xx}\|_{0,\omega} \|\psi_{xx}\|_{0,\omega},$$

so that (4.4) is a consequence of (4.6)-(4.9).

PROOF OF THEOREM 4.1: Let $f = g_{xx}$ be in $H_{\omega}^{-2}(I)$. Problem (4.1) is equivalent to the following:

$$(4.10) \quad \begin{cases} \text{Find } \psi \text{ in } V = H_{0,\omega}^2(I) \text{ such that, for any } \phi \text{ in } V: \\ \int_I \psi_{xx}(\phi\omega)_{xx} = \int_I g(\phi\omega)_{xx}. \end{cases}$$

The bilinear form a defined by: For any (χ, ϕ) in V^2 :

$$(4.11) \quad a(\phi, \chi) = \int_I \phi_{xx}(\chi\omega)_{xx},$$

is continuous and elliptic over V (see Lemma 4.2), and Lax-Milgram lemma gives the existence and uniqueness of a solution of (4.10) hence of (4.1).

Approximation of Problem 4.1

We are interested in approximating the solution of (4.1) by a polynomial of degree $\leq N$. We use a Galerkin method approach known as Spectral Method (see Gottlieb-Orszag [1] for more details); hence from (4.10) we derive an approximate problem:

$$(4.12) \quad \begin{cases} \text{Find } \psi_N \text{ in } V_N = S_N^2 \text{ such that, for any } \phi \text{ in } V_N: \\ \int_I \psi_{Nxx}(\phi\omega)_{xx} = \int_I g(\phi\omega)_{xx}. \end{cases}$$

From Lemma 4.2 we know that problem (4.12) is wellposed in the sense that there exists one and only one solution. Moreover, we derive from (4.10) and (4.12):

$$a(\psi - \psi_N, \phi) = 0 \quad \text{for any } \phi \text{ in } V_N,$$

so that (remind $\Pi_{2,N}^{2,0} \psi \in V_N$):

$$(4.13) \quad a(\psi - \psi_N, \psi - \psi_N) = a(\psi - \psi_N, \phi - \Pi_{2,N}^{2,0} \psi).$$

Due to Theorem 3.1 and Lemma 4.2, we then obtain the following

THEOREM 4.2: There exists one and only one solution ψ_N to problem (4.12); moreover it verifies, as soon as $\psi \in H_{\omega}^{\sigma}(I) \cap H_{0,\omega}^2(I)$:

$$(4.14) \quad \|\psi - \psi_N\|_{2,\omega} \leq C N^{2-\sigma} \|\psi\|_{\sigma,\omega}.$$

REMARK 4.1: The previous estimate is an optimal one in the sense that no polynomial of S_N^2 is asymptotically nearer from the solution ψ than the solution of the approximate problem.

REMARK 4.2: The previous theorem will be extended in a future paper where we shall consider a pseudospectral method (much more efficient from a computational point of view) for approximating a one-dimensional fourth order equation.

ACKNOWLEDGEMENT

The author would like to thank F. Murat for valuable discussion about interpolation results.

REFERENCES

- Canuto-Quarteroni [1] Approximation results for orthogonal polynomials in Sobolev spaces, Math. Comp., 38 (1982), pp. 67-86.
- [2] Variational methods in the theoretical analysis of spectral approximation, "Proc. of Symposium on Spectral Methods for Partial Differential Equations," R. G. Voigt, D. Gottlieb, and M. Y. Hussaini, eds., SIAM (Philadelphia) 1984, pp. 55-78.
- Gottlieb-Orszag [1] Numerical Analysis of Spectral Methods: Theory and Applications," CBMS Regional Conference Series in Applied Mathematics 26, SIAM (Philadelphia), 1977.
- Grisvard [1] Espaces intermediaires entre espaces de Sobolev avec poids, Ann. Scuola Norm. Sup. Pisa, 17 (1963).
- [2] Commutativité de 2 foncteurs d'interpolation et applications., J. Math. Pures Appl., 45 (1966), pp. 143-206
- Lions-Magenes [1] "Problemes aux limites non homogenes et applications," Dunod, 1968.
- Lions-Peetre [1] Sur une classe d'espaces d'interpolation, Publ. Math. a l'IHES, Paris, 19 (1964), pp. 5-68.

- Maday [1] Some spectral methods concerning a fourth order one-dimensional problem, to appear in Publications du Laboratoire d'Analyse Numérique de Paris VI (1984).
- Maday-Metivet [1] Estimation d'erreur pour l'approximation des equations de Stokes par une methode spectrale, La Rech. Aero., 5 (1983), pp. 237-244.
- [2] Chebyshev spectral approximation of Navier-Stokes equations in a two-dimensional domain, to appear in Publications du Laboratoire d'Analyse Numérique de Paris VI (1984).
- Maday-Quarteroni [1] Legendre and Chebyshev spectral approximation of Burger's equation, Numer. Math., 37 (1981), pp. 321-332.
- Nečas [1] "Les Methodes Directes dans la Theorie des Equations Elliptiques," Edition de l'Academie Tchecoslovaque des Sciences, Prague, 1967.
- Tadmor [1] The exponential accuracy of Fourier and Chebyshev differencing methods, ICASE Report No. 84-40, NASA NASA CR-172435, 1984.

1. Report No. NASA CR-172573 ICASE Report No. 85-17		2. Government Accession No.		3. Recipient's Catalog No.	
4. Title and Subtitle ANALYSIS OF SPECTRAL OPERATORS IN ONE-DIMENSIONAL DOMAINS				5. Report Date February 1985	
				6. Performing Organization Code	
7. Author(s) Yvon Maday				8. Performing Organization Report No. 85-17	
				10. Work Unit No.	
9. Performing Organization Name and Address Institute for Computer Applications in Science and Engineering Mail Stop 132C, NASA Langley Research Center Hampton, VA 23665				11. Contract or Grant No. NAS1-17070	
				13. Type of Report and Period Covered Contractor Report	
12. Sponsoring Agency Name and Address National Aeronautics and Space Administration Washington, D.C. 20546				14. Sponsoring Agency Code 505-31-83-01	
15. Supplementary Notes Langley Technical Monitor: J. C. South, Jr. Final Report					
16. Abstract We prove results concerning certain projection operators on the space of all polynomials of degree less than or equal to N with respect to a class of one-dimensional weighted Sobolev spaces. These results are useful in the theory of the approximation of partial differential equations with spectral methods.					
17. Key Words (Suggested by Author(s)) spectral methods interpolation spaces projection operators elliptic equations			18. Distribution Statement 64 - Numerical Analysis Unclassified - Unlimited		
19. Security Classif. (of this report) Unclassified	20. Security Classif. (of this page) Unclassified	21. No. of Pages 27	22. Price A03		

